



PROXIMATE COMPOSITION AND ORGANOLEPTIC PROPERTIES OF FISH CURED WITH TWO ENERGY SOURCES.

EBOCHUO, V.C*¹; EMUKA, E.C¹; ONYEME, O¹; EZENNADI, P.O¹ and AHAOTU, E.O²

¹*Department of Fisheries and Marine Technology, Imo State Polytechnic Umuagwo, Ohaji, Nigeria.* ²*Department of Animal Production Technology, Imo State Polytechnic Umuagwo, Ohaji, Nigeria.*

Abstract

The proximate composition and organoleptic properties of silver catfish (*Bagrus filamentosus*) cured with *Alchornea cordifolia* wood and Kainji Gas Klin (KGK) were determined.

Keywords

Proximate composition, Organoleptic properties, *Alchornea cordifolia* wood, *Bagrus filamentosus*, Kainji Gas Klin (KGK), cured fish.

INTRODUCTION

Fish is extremely perishable food item (Anyanwu, 2009 and Agbo *et al.*, 2002). The resultant effect is the easy decomposition of the fish at ambient state (Akinola *et al.*, 2006). Various factors are responsible for fish spoilage. Once fish is caught and upon comatose series of changes which include enzyme action (autolysis) chemical action (oxidative change) and microbial degradation begin to take place in the fish. This reason for fish processing is to give the product a form which is attractive to the consumers and to extend the storage life of fish. Davies (2005) suggested appropriate processing technologies to enable maximal use of raw

Fourteen pieces of proximate and cured with *Alchornea* *Bagrus filamentosus* organoleptic analysis. *cordifolia* wood had weighing between The result of the significantly ($P < 0.05$) 500g and 600g were proximate analysis higher lipid content purchased, killed, showed that fish cured than fish cured with deguted and washed to with Kainji Gas Klin Kainji Gas Klin remove blood and (KGK) had (18.71a \pm 0.08 and slime. The fish were significantly ($P < 0.05$) 10.10b \pm 0.10). The divided into two higher dry matter than result of the groups. Each group the same fish cured organoleptic analysis were immersed in ten with *Alchornea* showed that the liters of water that was *cordifolia* wood parameters taste and mixed with 300g of salt (92.40b \pm 0.10 and appearance for *Bagrus* to get 3% brine 90.00a \pm 0.10). Fish *filamentosus* cured solution and allowed cured with Kainji Gas with wood and gas to stand for an hour Klin (KGK) had were significantly respectively. The first significantly ($P < 0.05$) ($P < 0.05$) different group was cured with lesser moisture scores. However, *Alchornea cordifolia* content than same fish aroma, mouth feel and wood while the second cured with *Alchornea* general acceptability group was cured with *cordifolia* wood (8.90b so treated fish were Kainji Gas Klin (KGK) \pm 0.02 and not significantly for four hours 14.40a \pm 0.01). Fish different ($P > 0.05$). Cured respectively. Cured cured with *Alchornea* The two sources of samples were allowed *cordifolia* wood had energy could be used to cool at ambient insignificantly higher to cure fish since the temperature for 12 ($P > 0.05$) protein parameters examined hours and were content than Kainji Gas fall within the packed in a sterile Klin (KGK) cured fish recommended ranges. black polythene bag (59.98 \pm 0.09 and before being sent for 61.11b \pm 0.10). Fish

material and thus contribute to increase in economic profitability, The poor handling and preservation practice after capture affects the degree of spoilage of the fish

(Akinneye *et al.*, 2007). Fish is a major source of protein and its harvesting, handling remains sources of livelihood for millions of people as well as providing foreign exchange earning to many countries (Alyufaili and Opara, 2006).

Different types of processing methods include drying, smoking, freezing, chilling and brining but the most prominent fish preservation is smoke curing. This could be adduced to the fact that most of the fishing communities have no access to electricity in order to preserve through freezing their products. Electricity therefore is fast becoming a less reliable source of energy for fish processing and preservation (Akinola *et al.*, 2006). *Bagrus filamentosus* is a common commercial catch in Nigerian fresh waters especially in the Southern states of Nigeria. Most fish that cannot be sold as fresh fish are preserved by fishermen and fish mongers to extend their shelf-life. Through smoke curing, the energy sources used in this study are *Alchornea cordifolia* wood and Kainji gas kiln as control.

Alchornea cordifolia is a common member of the flora in Southern Nigeria. This plant is used for its medicinal uses, fodder and also for the supply of energy for cooking. Since gas is not within the reach of artisanal fishermen and fish mongers, *Alchornea cordifolia* could be utilised as a common energy source for smoking fish.

This study however determined the proximate composition of smoke cured *Bagrus filamentosus* processed with *Alchornea cordifolia* wood and natural gas and also the organoleptic properties of smoke cured *Bagrus filamentosus* processed with *Alchornea cordifolia fuel* wood and natural gas respectively.

MATERIALS AND METHODS

This study was conducted in the fish processing and preservation unit of the fisheries laboratory, Department of Fisheries and Marine Technology, Imo State Polytechnic Umuagwo, Nigeria.

EXPERIMENTAL MATERIALS

The experimental materials used include smoking kiln, Kainji gas kiln, machete, bowls, weighing balance, dissecting kits, gas, salt, water, fish (*Bagrus filamentosus*) and *Alchornea cordifolia fuel* wood.

COLLECTION AND PREPARATION OF FISH SAMPLES

Samples of *Bagrus filamentosus* were obtained from hospital water side market in Yenogoa, Bayelsa State, Nigeria. Fourteen (14) pieces of *Bagrus filamentosus* weighing between 400 grams and 600 grams were weighed on a manual weighing balance (Capacity brand: 20kg, Camry Emperors, made in China). Fish was killed, degut, washed and weighed. The fishes were divided into two groups. Each group was immersed in ten liters of water that was mixed with 300 grams for an hour before curing with *Alchornea cordifolia fuel* wood and gas. After smoking and gas curing, the fishes were weighed.

ORGANOLEPTIC ANALYSIS

Wood cured fish and gas cured fish were submitted to a ten man panel for sensory analysis using a seven point Hedonic scale to ascertain the following parameter:- taste, aroma, mouth feel, appearance and general acceptability according to the methods of Eyo (2001).

STATISTICAL ANALYSIS

The differences in proximate composition between smoke cured fish and gas cured fish was determined using student t- while the difference in organoleptic parameters between smoke cured fish and natural gas cured fish were determined by using Tukey test respectively.

BIOLOGICAL EVALUATION

The biological evaluation was done by obtaining the carcass weight of the fish. The fish samples were then cleaned and eviscerated. The weight after evisceration was then obtained as the dress weight. The weight of the cured fish was also obtained as weight after smoking.

RESULTS

Table 1. Weight Characteristics of *Bagrus filamentosus* cured with *Alchornea cordifolia fuel* wood and Kainji Gas Kiln

Samples	Live Weight	Dressed	Weighing	Total	% Weight
Weight	after Smoking	Weight Loss	Loss		
A	500	400	170	330	66.0
B	600	500	207	393	65.5

Average Weight Loss (%) = 65.7

Table 2. Proximate Composition of *Bagrus filamentosus* cured with *Alchornea cordifolia fuel* wood and Kainji Gas Kiln

T. S	% DM	%MO	%C.P	%LI	%Ash	% C.F	NFE
T ₁	90.00 ^a ± 0.10	14.40 ^a ±0.01	61.11 ^b ± 0.10	18.71 ^a ±0.08	11.50 ^a ±0.01	4.30 ^a ±0.02	2.30 ^a ±0.05
T ₂	92.40 ^b ±0.10	8.90 ^b ±0.02	59.98 ^a ±0.09	10.10 ^b ±0.10	5.40 ^b ±0.02	1.50 ^b ±0.03	2.56 ^a ±1.02

DM= Dry Matter; MO= Moisture; C.P= Crude protein; LI=Lipid; C.F= Crude fibre, NFE= Nitrogen free extract.

Where the same superscripts appear in the same column, there is significant different at (P<0.05).

T₁= *Alchornea cordifolia fuel* wood curing

T₂= Kainji Gas Kiln curing (KGK)

Table 3. Organoleptic Parameters of *Bagrus filamentosus* cured with *Alchornea cordifolia fuel* wood and Kainji Gas Kiln

Treatments	PARAMETERS				
	Taste	Appearance	Aroma	Mouth feel	General Acceptability
Wood	4.00±0.40 ^a	4.07±0.42 ^a	6.05±0.38 ^b	5.47±0.38 ^a	6.53±0.04 ^b
Gas	6.00±0.30 ^b	5.72±0.25 ^b	6.34±0.27 ^b	5.32±0.37 ^b	6.51±0.28 ^b

The study showed that *Bagrus filamentosus fish smoked with* cured with Kainji Gas Kiln had significantly (P<0.05) higher dry matter than the same fish cured with *Alchornea cordifolia fuel* wood as shown in table 1 above. Fish cured with Kainji Gas Kiln had significantly (P<0.05) lesser moisture than fish smoked with *Alchornea cordifolia fuel* wood as shown in table 2. The parameters' taste and appearance for *Bagrus filamentosus* cured with *Alchornea cordifolia fuel* wood and gas were significantly (P<0.05) different. However, aroma, mouth feel and general acceptability of the treated fish were not significantly different (P>0.05).

DISCUSSION

The average moisture loss (65.7%) from the cured *Alchornea cordifolia fuel* wood and Kainji Gas Kiln is similar to the value of less than 65.00% recommended by Bene and Heck (2005). The values recommended by Ebochuo and Oparaejiaku (2017) are e\within the range of 10-12% with corresponds with the value obtained from this study.

Results in crude protein showed that KGK produced a significantly lower product than the wood cured product. This confirms the findings of Eyo (2001) that different processing methods affect nutritional value of processed food. Emuka *et al.*, (2015) stated that oven drying, solar drying, smoking and boiling had different impacts on nutritional values of *S. Clarias Trachrus trecae* and *C. gariepinus* so treated. Emuka, *et al.*, (2015) further stated that processing methods which lowers the lipid values help to prevent oxidation and rancidity in fish preservation. Richardson and Goff (2001) advanced that the amount of heat from heat source significantly denatured the fibre in the muscle of peanid shrimps.

CONCLUSION

Alchornea cordifolia fuel wood which is abundant qualify for use in fish smoking. Both sources do not exceed of fall below the operating standard in Nigeria.

REFERENCES

- Agbon, A.O, Ezeri, G.N.O, Ikenwiewe, B.N, Alegbeleye, N.O and Akomolade, D.J (2002). A comparative study of different storage methods on the shelf –life of smoked current fish. J.Aquatic. Sci.. 17 (20: 134-136.
- Akinola, O.A, Akinfemi, A.A and Bolayi, B.O (2006). Evaluation of traditional and solar drying systems towards enhancing fish storage and preservation in Nigeria (Abeokuta Local Government Council:- A case study) J.Fish Int., 1: (2-4): 44-49.
- Akinneye, J.O, Amoo, I.A and Arannilewa, S.T (2007). Effect of drying methods on the nutritional composition of three species (*Bonga spp*; *Sardinella spp* and *Heterotis niloticus*). J. Fish Int., 2 (1): 99-103.
- Alyufaili, M.S and Opara, C.U (2006). Study of Fisheries post-harvest industry in the Sultanate of Oman. Part 1: Handling and Marketing System of Fresh Fish. J. Fish In; 1 (2-4): 144-149.
- Anyanwu, M.O (2009). Preservation of Smoked Fish from *Dermestes Maculatus* (Colcoptera dermestidae) Infestation Using *Piper guineensis* and *Xylopiia aethiopica*. Student Project. Department of Fishery and Marine Technology, Imo State Polytechnic Umuagwo, Nigeria. 17 pp.

- Bene, C. and Heck . S. (2005). Fisheries and the millennium development goals. Solution for Africa. NAGA 28: 8 – 13.
- Davies, R.M (2005). Development of Appropriate Technology of Fish Processing in Nigeria. A paper presented at a one day workshop on Intensive Fish Farming.
- Ebochuo, V.C and Oparaejiaku, J (2017). Nutritive value and organoleptic attributes of *Xylopia aethiopica* pre-treated smoke cured *Clarias gariepinus* (Burchell, 1822). Direct Res. J. Biol. Biotechnol. Sci. 3 (3): 56 – 59.V
- Emuka, E.C; Onyeme, O and Ezennadi, P.O (2015). Proximate composition and Organoleptic Properties of Fish cured with two Energy Sources. Student Project. Department of Fisheries and Marine Technology, Imo State Polytechnic Umuagwo, Nigeria. 39pp.
- Eyo, A.A (2001). Fish Processing Technology in the Tropics. NIFFR Press, New Bussa, Nigeria.
- Richardson, A.S and Goff, E.R (2001). Effects of Temperature and intraspecific interaction on the development of *Dermestes maculatus* (Coleoptera; Dermestidae). Journal of Medical Entomology 38: 347-351